
# MME 2509 Materials Processing Laboratory

## Metal Oxide Synthesis via Organic Precursor Method


Assist. Prof. Dr. Çınar Öncel

 $Sr_4Al_2O_7$ ,  $Sr_3Al_2O_6$ ,  $SrAl_2O_4$ ,  $SrAl_4O_7$ ,  $SrAl_{12}O_{19}$ 





### **Organic Precursor Method**

- Involves dissolution of cation salts in a proper liquid
- Metal cations are distributed homogeneous in molecular level.
- Organic precursor as chelating agents dissolves in the same solution.
- Hydroxyl or carboxyl ends capture positive metal ions until decomposition











Citric acid



Ethylene diaminetetraacetic acid (EDTA).

J:N=C O  $0 \ge 0^{-Al} = 0 \ge 0^{-Al} = 0$  $\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}_{3} \begin{bmatrix} AI^{3+} \end{bmatrix} Mn^{2+} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}_{2}$ 



hydrogen	9 ( <del>)</del>		1070		1070	10		15		1.5653	2.20	10.000		0.575	8.8%	10.0	4.0 S	helium
Η Ĥ Ι																		He
1.0079																		4.0026
lithium 3	beryllium <b>4</b>												boron 5	carbon 6	nitrogen 7	oxygen 8	fluorine 9	neon 10
Li	Be												B	C	Ň	0	F	Ne
6.941	9.0122												10.811	12.011	14.007	15.999	18.998	20.180
sodium 11	magnesium 12												aluminium 13	silicon 14	phosphorus 15	sulfur 16	chlorine 17	argon 18
122.000	Statistics of the second second													12334/82		2020	2012	123323
Na	Mg												AI	Si	Ρ	S	CI	Ar
22.990 potassium	24.305 calcium		scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	26.982 gallium	28.086 germanium	30.974 arsenic	32.065 selenium	35.453 bromine	39.948 krypton
19	20		21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca		Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
39.098	40.078		44.956	47.867	50.942	51.996	54.938	55.845	58.933	58.693	63.546	65.39	69.723	72.61	74.922	78.96	79.904	83.80
rubidium 37	strontium 38		yttrium <b>39</b>	zirconium 40	niobium 41	molybdenum 42	technetium 43	ruthenium 44	rhodium 45	palladium <b>46</b>	silver 47	cadmium 48	indium 49	tin 50	antimony 51	tellurium 52	iodine 53	xenon 54
Rb	Sr		Y	Zr	Nb	Мо	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Те		Xe
85.468	87.62		88,906	91.224	92.906	95.94	[98]	101.07	102.91	106.42	107.87	112.41	114.82	118.71	121.76	127.60	126.90	131.29
caesium 55	barium 56	57-70	lutetium 71	hafnium <b>72</b>	tantalum 73	tungsten 74	rhenium 75	osmium 76	iridium 77	platinum <b>78</b>	gold 79	mercury 80	thallium 81	lead 82	bismuth 83	polonium 84	astatine 85	radon 86
Construction of the second	Contraction of the second s	*	12000			Ŵ	- Parodinin			and the second second		1000 C 1000 C 20	TI		100000	1	and the second	17.5 N. 5 N. 5 N.
Cs	Ba	$\mathbf{x}$	Lu	Hf	Та		Re	Os	Ir	Pt	Au	Hg		Pb	Bi	Po	At	Rn
132.91 francium	137.33 radium		174.97 lawrencium	178.49 rutherfordium	180.95 dubnium	183.84 seaborgium	186.21 bohrium	190.23 hassium	192.22 meitnerium	195.08 ununnilium	196.97 unununium	200.59 ununbium	204.38	207.2 ununquadium	208.98	209	[210]	[222]
87	88	89-102	103	104	105	106	107	108	109	110	111	112		114				
Fr	Ra	* *	Lr	Rf	Db	Sg	Bh	Hs	Mt	Uun	Uuu	Uub		Uuq				
[223]	[226]		[262]	[261]	[262]	[266]	[264]	[269]	[268]	[271]	[272]	2771		[289]				

*Lanthanide series	lanthanum 57	58	praseodymium 59	60	61	samarium 62	europium 63	gadolinium <b>64</b>	terbium 65	dysprosium 66	holmium 67	erbium 68	thulium 69	ytterbium 70
~~~	La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb
	138.91	140.12	140.91	144.24	[145]	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.04
* * Actinide series	actinium 89	thorium 90	protactinium 91	uranium 92	neptunium 93	plutonium <b>94</b>	americium 95	curium 96	berkelium 97	californium 98	einsteinium 99	fermium 100	mendelevium 101	nobelium 102
	Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No
	[227]	232.04	231.04	238.03	[237]	[244]	[243]	[247]	[247]	[251]	[252]	[257]	[258]	[259]

Oxide A	Oxide B	Oxide C
Sr <sub>3</sub> Al <sub>2</sub> O <sub>6</sub>	3%Cr-doped Al <sub>2</sub> O <sub>3</sub>	SrAl <sub>2</sub> O <sub>4</sub>

Sources: Sr(NO<sub>3</sub>)<sub>2</sub>, Al(NO<sub>3</sub>)<sub>3</sub>, Chromium Nitrate, citric acid monohydrate

You will calculate how much sources you need to produce 1 gr of powder  $(Sr_3Al_2O_6, SrAl_2O_4 \text{ or } 3\%Cr-doped Al_2O_3)$  and citric acid monohydrate \* You must use 1 citric acid molecule for each cation.



~300<sup>0</sup>C

### 800<sup>o</sup>C – 30 minutes

### → 1100°C – 1 hour







\*\*\*All in air

